



ÚOCHB AV
IOCB PRAGUE


Ústav organické chemie a biochemie
Akademie věd České republiky, v. v. i.
Institute of Organic Chemistry and Biochemistry
of the Czech Academy of Sciences

PRESS RELEASE

How to easily and inexpensively produce fluorescent nanoparticles for medical applications in a nuclear reactor

Prague, November 9, 2018 – Under the leadership of Petr Cígler from the Institute of Organic Chemistry and Biochemistry and Martin Hrubý from the Institute of Macromolecular Chemistry, both of which are part of the Czech Academy of Sciences, a team of researchers has developed a revolutionary method for the easy and inexpensive production of irradiated nanodiamonds and other nanomaterials suitable for use in highly sensitive diagnostics of diseases, including various types of cancer. Their article was recently published in the scientific journal *Nature Communications*.

Diagnosing diseases and understanding the processes that take place within cells at the molecular level require sensitive and selective diagnostic instruments. Today, scientists can monitor magnetic and electric fields in cells at a resolution of several dozen nanometers and with remarkable sensitivity thanks to crystal defects in the particles of certain inorganic materials. A nearly ideal material for these purposes is diamond. Compared with the diamonds used in jewelry, the ones intended for applications in diagnostics and nanomedicine – nanodiamonds – are approximately a million times smaller and are produced synthetically from graphite at high pressure and temperatures.



A pure nanodiamond, though, doesn't reveal much about its environment. First, its crystal lattice must be damaged under controlled conditions to create special defects, so-called nitrogen-vacancy centers, which enable optical imaging. The damage is most commonly created by irradiating nanodiamonds with fast ions in particle accelerators. These accelerated ions are capable of knocking carbon atoms out of the crystal lattice of a nanodiamond, leaving behind holes known as vacancies, which at high temperatures then pair with nitrogen atoms present in the crystal as contaminants. The newly formed nitrogen-vacancy centers are a source of fluorescence, which can then be observed. It's precisely this fluorescence that gives nanodiamonds immense potential for applications in medicine and technology. A fundamental restriction to the use of these materials on a broader scale, however, is the great cost and poor efficiency of irradiating ions in an accelerator, which prevents the generation of this exceptionally valuable material in larger quantities.

The team of scientists from several research centers headed by Petr Cígler and Martin Hrubý has recently published an article in the journal *Nature Communications* describing an entirely new method of irradiating nanocrystals. In place of costly and time-consuming irradiation in an accelerator, the scientists exploited irradiation in a nuclear reactor, which is much faster and far less expensive.

But it wasn't quite that simple. The scientists had to employ a trick – in the reactor, neutron irradiation splits boron atoms into very light and fast ions of helium and lithium. The nanocrystals must first be dispersed in molten boron oxide and then subjected to neutron irradiation in a nuclear reactor. Neutron capture by boron nuclei produces a dense shower of helium and lithium ions, which have the same effect within the nanocrystals as the ions produced in an accelerator: the controlled creation of crystal defects. The high density of this particle shower and the use of a reactor to irradiate a much larger quantity of material mean that it is easier and far more affordable to produce dozens of grams of rare nanomaterial at once, which is approximately one thousand times more than scientists have thus far been able to obtain through comparable irradiation in accelerators. The method has proven successful not only in the creation of defects in the lattice of nanodiamonds but of another nanomaterial as well – silicon carbide. For this reason, scientists hypothesize that the method could find universal application in the large-scale production of nanoparticles with defined defects.

The new method utilizes the principle applied in boron neutron capture therapy (BNCT), in which patients are administered a boron compound. Once the compound has collected in the tumor, the patient receives radiation therapy with neutrons, which split the boron nuclei into ions of helium and lithium. These then destroy the tumor cells that the boron has collected in. This principle taken from experimental cancer treatment thus has opened the door to the efficient production of nanomaterials with exceptional potential for applications in, among other areas, cancer diagnostics.

Article: Jan Havlík, Vladimíra Petráková, Jan Kučka, Helena Raabová, Dalibor Pánek, Václav Štěpán, Zuzana Zlámalová Cílová, Philipp Reineck, Jan Štursa, Jan Kučera, Martin Hrubý a Petr Cígler: Extremely rapid isotropic irradiation of nanoparticles with ions generated in situ by a nuclear reaction. *Nature Communications* **2018**, 9, 4467. DOI: [10.1038/s41467-018-06789-8](https://doi.org/10.1038/s41467-018-06789-8).

The Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences / IOCB Prague (www.iocb.cz) is a leading internationally recognized scientific institution whose primary mission is the pursuit of basic research in chemical biology and medicinal chemistry, organic and materials chemistry, chemistry of natural substances, biochemistry and molecular biology, physical chemistry, theoretical chemistry, and analytical chemistry. An integral part of the IOCB Prague's mission is the implementation of the results of basic research in practice. Emphasis on interdisciplinary research gives rise to a wide range of applications in medicine, pharmacy, and other fields.

--- END PRESS RELEASE ---

MEDIA RELATIONS:

Dušan Brinzanik (IOCB Prague – Communications): dušan.brinzanik@uochb.cas.cz,
mob: +420 731 609 271